

COMPOSITION

CEVAC® TREMOR L contains the Calnek strain of Avian Encephalomyelitis virus in live, freeze dried form. The embryonated hen eggs used in the production of the vaccine are obtained from specified-pathogen-free (SPF) flocks.

INDICATIONS

For the active immunisation of healthy chickens against Avian Encephalomyelitis. The progeny of hens vaccinated according to the recommended program will have maternally derived antibodies protecting it from Avian Encephalomyelitis.

CONTRA-INDICATIONS

CEVAC® TREMOR L should not be used during lay as it will cause a drop in production and the vaccine strain will pass on to the eggs for a period a short period after vaccination.

ADMINISTRATION AND DOSAGE

Pullets should be vaccinated with CEVAC® TREMOR L after 10 weeks of age and 4 weeks before the onset of laying by drinking water or wing web administration. Wing Web: Insert the applicator into the webbed portion of the wing avoiding feathers, muscle, bone and blood vessels (0.01 ml dose/bird). Vaccine should be used within one hour.

STORAGE

- Store vaccine between +2°C and +8°C or 35°F and 45°F
- Protect from light

PRESENTATIONS

1,000 dose vials 20 x 1,000 dose vials / box

WARNING

To prevent losses in egg production and/or vertical transmission of avian encephalomyelitis virus to progeny, DO NOT ADMINISTER THIS VACCINE TO CHICKENS DURING OR NEAR EGG PRODUCTION!