

COCCI - VAC

Coccidiosis vaccine (Live Oocysts)

DISCRIPTION AND INDICATIONS

COCCI-VAC is used for the immunization of poultry against coccidiosis. It contains live oocysts. Immunity develops in 10 to 14 days after vaccination.

ADMINISTRATION

Method of dilution

Age of chicken	Vaccine	Water for dilution
4 to 28 day	1,000 doses	10ℓ

VACCINATION PROGRAM

Vaccinate birds 4 to 14 days old (preferably 7 ~ 10 days).

PRECAUTIONS

1. Vaccinate only healthy fowl.
2. Discontinue the use of any medicated drinking water or disinfectants 24hours before and after vaccination.
3. Do not use on chickens receiving feed containing a coccidiostat or high levels of antibiotics.

4 . Virus vaccine may be given at the same time in the same drinking water. Always dilute COCCI-VAC with full amount of water before adding virus vaccine.

5. Put stabilizer into the water and mix well for 2 to 3 hours.

6. Use entire contents of each vial after opening.

7. Incinerate the vaccine container and any unused contents.

8. Do not use frozen vaccine.

STORAGE

Store at 2°to 7°C and protect from light. Do not freeze.

PACKING SIZE

1,000 doses X 10 vials with stabilizer.

- COMPOSITION/PACKAGING MATERIALS

1. COMPOSITION

I. Name :

Coccidium vaccine (Modified Live Oocysts.)

II. Appearance;

Liquid of a yellow-brownish color.

III. Composition of Vaccine (1,000 doses / vial)

Species of oocyst	No.of oocyst
<i>E. necatrix</i>	200,000
<i>E. tenella</i>	150,000
<i>E. hagani</i>	800,000
<i>E. acervulina</i>	800,000
<i>E. maxima</i>	400,000
<i>E. brunetti</i>	800,000
<i>E. praecox</i>	800,000
<i>E. mivati</i>	800,000
* Storage solution.....	4.0 Mℓ
* Storage solution	
Potassium dichromate.....	25g
Distilled water.....	1,000 Mℓ