Volvac® AC Plus Emul Bacterin

Coriza Infecciosa ABC/ Infectious Coryza

Reg. SAGARPA B-0171-123

FORMULA:

Trivalent bacterin with A, B, C Serogroups of *Avibacterium paragallinarum*, inactivated and oil emulsified. A volume of 0.5ml contains a minimum titer of 10^{8.0} CFU/dose or 10^{8.0} EID 50/dose for each serogroup

INDICATIONS:

The antigenicity of *Avibacterium paragallinarum* strains and A, B, C combined serogroups induce a solid and long lasting immunity. These antigens, have been adjuvated and oil emulsified are recommended for the first and booster revaccination. For the immunization of healthy birds against Avian Infectious Coryza.

DOSE AND ADMINISTRATION ROUTE:


0.5 ml per bird.

Subcutaneous, in third half of the back of the neck.

- Store out of direct sunlight, in a refrigerator between 2° to 7°C.
- I Do not freeze.
- I Do not keep remains for subsequent vaccinations.
- Incinerate all empty bottles after vaccination.
- Read carefully the instructions written on the label before use.
- Keep out of children's reach.
- I Consult your poultry veterinarian.

PRESENTATIONS:

Plastic Bottles 500 ml/1,000 doses





Productos / Aves / Vacunas

Abrir Version para Impresión

© 2007 Esta información es propiedad de Boehringer Ingelheim Vetmedica, S.A. de C.V., cualquier reproducción parcial o total deberá hacerse con la autorización de la Compañía. Tel.: 52 (33) 3668-8000

Arriba