OVO-4

Adjuvanted inactivated vaccine against Infectious Bronchitis, Newcastle Disease, Egg Drop Syndrome, and Swollen Head Syndrome INJECTABLE EMULSION

COMPOSITION

Each dose of vaccine contains:

- Inactivated Infectious Bronchitis Virus, titrating prior to inactivation minimum 10^{6.7} FID50
- Inactivated Newcastle Disease Virus, titrating prior to inactivation minimum 108.0 EID50
- Inactivated Egg Drop Syndrome Virus, titrating prior to inactivation minimum $10^{6.0}$ CCID50
- Inactivated Swollen Head Syndrome Virus, titrating prior to inactivation minimum $10^{5.0}$ CCID50
- Thiomersal, maximum 0.015mg
- Oil excipient, q.s. 0.3mL

INDICATION

Active immunization of future laying pullets and future breeding pullets against Infectious Bronchitis, Newcastle Disease, Egg Drop Syndrome, and Swollen Head Syndrome, as a booster for the respective live vaccines.

ADMINISTRATION AND DOSAGE

- Subcutaneous and/or intramascular route
- Dosage: one 0.3mL dose per bird

One single injection 2 to 4 weeks prior to the laying season

PRECAUTIONS

- Shake well prior to use
- Vaccinate only healthy animals
- Apply usual aseptic procedures
- Do not use syringes with natural rubber-based or butyl-derived elastomer pistons

WITHDRAWAL PERIOD

Not applicable

STORAGE

Store between +2°C and +8°C, protected from light

FOR VETERINARY USE ONLY